

# HELICENES: CHIRAL PLATFORMS WITH FASCINATING PROPERTIES






## Jeanne Crassous

CNRS - Univ. Rennes,  
Institut des Sciences Chimiques de Rennes  
ISCR - UMR 6226, F-35000 Rennes, France

E-mail: [jeanne.crassous@univ-rennes1.fr](mailto:jeanne.crassous@univ-rennes1.fr)

*Ortho*-fused aromatic rings form helically shaped chiral molecules such as carbo[6]helicenes, that wind in a left-handed (*M*) or a right-handed (*P*) sense.<sup>1</sup> The helical topology combined with extended  $\pi$ -conjugation provides helicenes with peculiar properties such as strong photophysical and chiroptical properties (high optical rotation values, intense electronic circular dichroism and circularly polarized emission). The molecular engineering of helicenes using organometallic and heteroaromatic chemistries offers a convenient way to tune the properties of these helically shaped  $\pi$ -ligands. Indeed, their combination with metallic or organic assembling units leads to chiral materials with appealing properties (circularly polarized phosphorescence, magnetochirality, spin selectivity) for applications in materials science (Circularly Polarized OLEDs, Chiroptical Switches, Spintronics). I will present a set of representative examples.<sup>2</sup>



## References

1) a) C. -F. Chen, Y. Shen, *Helicenes Chemistry: From Synthesis to Applications*. Springer, Berlin, **2017**; b) M. Gingras, *Chem. Soc. Rev.* **2013**, *42*, 1051; c) K. Dhbaibi, L. Favereau, J. Crassous *Chem. Rev.* **2019**, *119*, 8846; d) J. Crassous, I. G. Stará, I. Starý (Eds) *Helicenes - Synthesis, Properties and Applications*. Wiley, **2022**.

2) a) E. S. Gauthier, L. Abella, N. Hellou, B. Darquié, E. Caytan, T. Roisnel, N. Vanthuyne, L. Favereau, M. Srebro-Hooper, J. A. G. Williams, J. Autschbach, J. Crassous, *Angew. Chem. Int. Ed.* **2020**, *59*, 8394; b) K. Dhbaibi, L. Abella, S. Meunier-Della-Gatta, T. Roisnel, N. Vanthuyne, B. Jamoussi, G. Pieters, B. Racine, E. Quesnel, J. Autschbach, J. Crassous, L. Favereau, *Chem. Sci.* **2021**, *12*, 5522; c) M. Atzori, K. Dhbaibi, H. Douib, M. Grasser, V. Dorcet, I. Breslavetz, K. Paillet, O. Cador, G. L. J. A. Rikken, B. Le Guennic, J. Crassous, F. Pointillart, C. Train, *J. Am. Chem. Soc.* **2021**, *143*, 2671; d) R. Rodríguez, C. Naranjo, A. Kumar, P. Matozzo, T.-K. Das, Q. Zhu, N. Vanthuyne, R. Gómez, R. Naaman, L. Sánchez, J. Crassous, *J. Am. Chem. Soc.* **2022**, *144*, 7709; e) K. Dhbaibi, M. Grasser, H. Douib, V. Dorcet, O. Cador, N. Vanthuyne, F. Riobé, O. Maury, S. Guy, Amina Bensalah-Ledoux, B. Baguenard, Geert L. J. A. Rikken, C. Train, B. Le Guennic, M. Atzori, F. Pointillart, J. Crassous, *Angew. Chem. Int. Ed.* **2023**, *62*, e202215558.